



TITLE:

THE VANISHING CYCLES OF TYPES
 $A_{\frac{1}{2}\infty}$ AND
 $D_{\frac{1}{2}\infty}$ (Geometry on Real
Closed Field and its Application to
Singularity Theory)

AUTHOR(S):

SAITO, KYOJI

CITATION:

SAITO, KYOJI. THE VANISHING CYCLES OF TYPES $A_{\frac{1}{2}\infty}$ AND $D_{\frac{1}{2}\infty}$ (Geometry on Real Closed Field and its Application to Singularity Theory). 数理解析研究所講究録 2011, 1764: 70-70

ISSUE DATE:

2011-09

URL:

<http://hdl.handle.net/2433/171399>

RIGHT:

THE VANISHING CYCLES OF TYPES $A_{\frac{1}{2}\infty}$ AND $D_{\frac{1}{2}\infty}$

KYOJI SAITO

ABSTRACT. We introduce two real entire functions $f_{A_{\frac{1}{2}\infty}}$ and $f_{D_{\frac{1}{2}\infty}}$ in two variables, having only two critical values 0 and 1. Associated maps $\mathbf{C}^2 \rightarrow \mathbf{C}$ define topologically locally trivial fibrations over $\mathbf{C} \setminus \{0, 1\}$. The critical points over 0 and 1 are ordinary double points, whose associated vanishing cycles in the generic fiber span its middle homology group and their intersection diagram forms the bi-partite decomposition of quivers of type $A_{\frac{1}{2}\infty}$ and $D_{\frac{1}{2}\infty}$, respectively. Coxeter element of type $A_{\frac{1}{2}\infty}$ and $D_{\frac{1}{2}\infty}$ are introduced as the product of the monodromies of the fibrations around 0 and 1. We describe the spectra of the intersection form (normalized in the interval $[0, 4]$) and the Coxeter elements (normalized in the interval $(-\frac{1}{2}, \frac{1}{2})$).

The present note is taken from the abstract of a preprint of the author: RIMS-1710 (Jan. 2011), Coxeter elements for vanishing cycles of types $A_{\frac{1}{2}\infty}$ and $D_{\frac{1}{2}\infty}$, which is going to appear elsewhere.